


SEP 25 2006

Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01

AMENDMENTS TO THE CLAIMS

1. (Currently Amended) A compound represented by the structural formula (I):



or pharmaceutically acceptable isomers, salts, solvates or esters of the compound of Formula (I), wherein in Formula (I) above:

X, Y and Z can be the same or different and each is independently selected from the group consisting of  $-\text{CH}_2-$ ,  $-\text{CH}(\text{alkyl})-$  and  $-\text{C}(\text{alkyl})_2-$ ;

$\text{Q}^1$  and  $\text{Q}^2$  can be the same or different and each is independently selected from the group consisting of H, -G,  $-(\text{C}_1-\text{C}_{30} \text{ alkylene})-\text{G}$ ,  $-\text{OR}^6$ ,  $-\text{OC}(\text{O})\text{R}^6$ ,  $-\text{OC}(\text{O})\text{OR}^9$ ,  $-\text{OC}(\text{O})\text{NR}^6\text{R}^7$ , and  $-\text{L}-\text{M}$ ;

$\text{Q}^3$  is 1 to 5 substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, -G,  $-(\text{C}_1-\text{C}_{30} \text{ alkylene})-\text{G}$ ,  $-\text{OR}^6$ ,  $-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{OR}^6$ ,  $-\text{C}(\text{O})\text{R}^6$ ,  $-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{C}(\text{O})\text{R}^6$ ,  $-\text{C}(\text{O})\text{OR}^6$ ,  $-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{C}(\text{O})\text{OR}^6$ ,  $-\text{OC}(\text{O})\text{R}^6$ ,  $-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{OC}(\text{O})\text{R}^6$ ,  $-\text{OC}(\text{O})\text{OR}^9$ ,  $-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{OC}(\text{O})\text{OR}^9$ ,  $-\text{CH}=\text{CH}-\text{C}(\text{O})\text{R}^6$ ,  $-\text{CH}=\text{CH}-\text{C}(\text{O})\text{OR}^6$ ,  $-\text{C}\equiv\text{C}-\text{C}(\text{O})\text{OR}^6$ ,  $-\text{C}\equiv\text{C}-\text{C}(\text{O})\text{R}^6$ ,  $-\text{O}-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{OR}^6$ ,  $-\text{O}-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{C}(\text{O})\text{R}^6$ ,  $-\text{O}-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{C}(\text{O})\text{OR}^6$ , -CN,  $-\text{O}-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{C}(\text{O})\text{NR}^6\text{R}^7$ ,  $-\text{O}-\text{C}(\text{O})\text{NR}^6\text{NR}^7\text{C}(\text{O})\text{OR}^6$ ,  $-\text{O}-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{C}(\text{O})\text{NR}^6\text{NR}^7\text{C}(\text{O})\text{OR}^6$ ,  $-\text{O}-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{C}(\text{O})(\text{aryl})-\text{N}_3$ ,  $-\text{OC}(\text{O})-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{C}(\text{O})\text{OR}^6$ ,  $-\text{C}(\text{O})\text{NR}^6\text{R}^7$ ,  $-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{C}(\text{O})\text{NR}^6\text{R}^7$ ,  $-\text{OC}(\text{O})\text{NR}^6\text{R}^7$ ,  $-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{OC}(\text{O})\text{NR}^6\text{R}^7$ ,  $-\text{NO}_2$ ,  $-\text{NR}^6\text{R}^7$ ,  $-(\text{C}_1-\text{C}_{10} \text{ alkylene})-\text{NR}^6\text{R}^7$ ,

Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01

-O-(C<sub>2</sub>-C<sub>10</sub> alkylene)-NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>C(O)R<sup>7</sup>, -NR<sup>6</sup>C(O)OR<sup>9</sup>,  
-NR<sup>6</sup>C(O)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>S(O)<sub>0-2</sub>R<sup>9</sup>, -N(S(O)<sub>0-2</sub>R<sup>9</sup>)<sub>2</sub>, -CHNOR<sup>6</sup>, -C(O)NR<sup>6</sup>R<sup>7</sup>,  
-C(O)NR<sup>6</sup>NR<sup>6</sup>R<sup>7</sup>, -S(O)<sub>0-2</sub>NR<sup>6</sup>R<sup>7</sup>, -S(O)<sub>0-2</sub>R<sup>9</sup>, -O-C(O)-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)NR<sup>6</sup>R<sup>7</sup>,  
-OC(O)-(C<sub>1</sub>-C<sub>10</sub> alkylene)-NR<sup>6</sup>C(O)O-(alkylaryl), -P(O)(OR<sup>10</sup>)<sub>2</sub>,  
-(C<sub>1</sub>-C<sub>10</sub> alkylene)-OSi(alkyl)<sub>3</sub>, -CF<sub>3</sub>, -OCF<sub>3</sub>, halo, alkoxyalkoxy, alkoxyalkoxyalkoxy,  
alkoxycarbonylalkoxy, alkoxyarylalkoxy, alkoxyiminoalkyl, alkylidoyl, allyloxy, aryl, arylalkyl,  
aryloxy, arylalkoxy, aroyl, aroyloxy, aroylaroyloxy, arylalkoxycarbonyl, benzoylbenzoyloxy,  
heteroaryl, heteroarylalkyl, heteroarylalkoxy, dioxolanyl, heterocyclyl, heterocyclylalkyl,  
heterocyclylcarbonyl, heterocyclylcarbonylalkoxy and -L-M;

Q<sup>4</sup> is 1 to 5 substituents independently selected from the group consisting of alkyl,  
alkenyl, alkynyl, -G, -(C<sub>1</sub>-C<sub>30</sub> alkylene)-G, -OR<sup>6</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-OR<sup>6</sup>, -C(O)R<sup>6</sup>,  
-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)R<sup>6</sup>, -C(O)OR<sup>6</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)OR<sup>6</sup>, -OC(O)R<sup>6</sup>,  
-(C<sub>1</sub>-C<sub>10</sub> alkylene)-OC(O)R<sup>6</sup>, -OC(O)OR<sup>9</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-OC(O)OR<sup>9</sup>, -CH=CH-C(O)R<sup>6</sup>,  
-CH=CH-C(O)OR<sup>6</sup>, -C≡C-C(O)OR<sup>6</sup>, -C≡C-C(O)R<sup>6</sup>, -O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-OR<sup>6</sup>,  
-O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)R<sup>6</sup>, -O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)OR<sup>6</sup>, -CN,  
-O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)NR<sup>6</sup>R<sup>7</sup>, -O-C(O)NR<sup>6</sup>NR<sup>7</sup>C(O)OR<sup>6</sup>, -O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-  
C(O)NR<sup>6</sup>NR<sup>7</sup>C(O)OR<sup>6</sup>, -O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)(aryl)-N<sub>3</sub>, -OC(O)-(C<sub>1</sub>-C<sub>10</sub> alkylene)-  
C(O)OR<sup>6</sup>, -C(O)NR<sup>6</sup>R<sup>7</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)NR<sup>6</sup>R<sup>7</sup>,  
-OC(O)NR<sup>6</sup>R<sup>7</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-OC(O)NR<sup>6</sup>R<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>6</sup>R<sup>7</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-NR<sup>6</sup>R<sup>7</sup>,  
-O-(C<sub>2</sub>-C<sub>10</sub> alkylene)-NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>C(O)R<sup>7</sup>, -NR<sup>6</sup>C(O)OR<sup>9</sup>,  
-NR<sup>6</sup>C(O)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>S(O)<sub>0-2</sub>R<sup>9</sup>, -N(S(O)<sub>0-2</sub>R<sup>9</sup>)<sub>2</sub>, -CHNOR<sup>6</sup>, -C(O)NR<sup>6</sup>R<sup>7</sup>,  
-C(O)NR<sup>6</sup>NR<sup>6</sup>R<sup>7</sup>, -S(O)<sub>0-2</sub>NR<sup>6</sup>R<sup>7</sup>, -S(O)<sub>0-2</sub>R<sup>9</sup>, -O-C(O)-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)NR<sup>6</sup>R<sup>7</sup>,  
-OC(O)-(C<sub>1</sub>-C<sub>10</sub> alkylene)-NR<sup>6</sup>C(O)O-(alkylaryl), -P(O)(OR<sup>10</sup>)<sub>2</sub>,  
-(C<sub>1</sub>-C<sub>10</sub> alkylene)-OSi(alkyl)<sub>3</sub>, -CF<sub>3</sub>, -OCF<sub>3</sub>, halo, alkoxyalkoxy, alkoxyalkoxyalkoxy,  
alkoxycarbonylalkoxy, alkoxyarylalkoxy, alkoxyiminoalkyl, alkylidoyl, allyloxy, aryl, arylalkyl,  
aryloxy, arylalkoxy, aroyl, aroyloxy, aroylaroyloxy, arylalkoxycarbonyl, benzoylbenzoyloxy,

Reply under 37 CFR 1.116  
 Expedited Procedure  
 Technology Center 1624  
 Attorney Docket No. CV06025US01

heteroaryl, heteroarylalkyl, heteroarylalkoxy, dioxolanyl, heterocyclyl, heterocyclylalkyl, heterocyclylcarbonyl, heterocyclylcarbonylalkoxy and -L-M;

$Q^5$  is 1 to 5 substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, -G, -(C<sub>1</sub>-C<sub>30</sub> alkylene)-G, -OR<sup>6</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-OR<sup>6</sup>, -C(O)R<sup>6</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)R<sup>6</sup>, -C(O)OR<sup>6</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)OR<sup>6</sup>, -OC(O)R<sup>6</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-OC(O)R<sup>6</sup>, -OC(O)OR<sup>9</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-OC(O)OR<sup>9</sup>, -CH=CH-C(O)R<sup>6</sup>, -CH=CH-C(O)OR<sup>6</sup>, -C≡C-C(O)OR<sup>6</sup>, -C≡C-C(O)R<sup>6</sup>, -O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-OR<sup>6</sup>, -O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)R<sup>6</sup>, -O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)OR<sup>6</sup>, -CN, -O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)NR<sup>6</sup>R<sup>7</sup>, -O-C(O)NR<sup>6</sup>NR<sup>7</sup>C(O)OR<sup>6</sup>, -O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)NR<sup>6</sup>NR<sup>7</sup>C(O)OR<sup>6</sup>, -O-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)(aryl)-N<sub>3</sub>, -OC(O)-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)OR<sup>6</sup>, -C(O)NR<sup>6</sup>R<sup>7</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)NR<sup>6</sup>R<sup>7</sup>, -OC(O)NR<sup>6</sup>R<sup>7</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-OC(O)NR<sup>6</sup>R<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>6</sup>R<sup>7</sup>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-NR<sup>6</sup>R<sup>7</sup>, -O-(C<sub>2</sub>-C<sub>10</sub> alkylene)-NR<sup>6</sup>R<sup>7</sup>, -NR<sup>6</sup>C(O)R<sup>7</sup>, -NR<sup>6</sup>C(O)OR<sup>9</sup>, -NR<sup>6</sup>C(O)NR<sup>7</sup>R<sup>8</sup>, -NR<sup>6</sup>S(O)<sub>0-2</sub>R<sup>9</sup>, -N(S(O)<sub>0-2</sub>R<sup>9</sup>)<sub>2</sub>, -CHNOR<sup>6</sup>, -C(O)NR<sup>6</sup>R<sup>7</sup>, -C(O)NR<sup>6</sup>NR<sup>6</sup>R<sup>7</sup>, -S(O)<sub>0-2</sub>NR<sup>6</sup>R<sup>7</sup>, -S(O)<sub>0-2</sub>R<sup>9</sup>, -O-C(O)-(C<sub>1</sub>-C<sub>10</sub> alkylene)-C(O)NR<sup>6</sup>R<sup>7</sup>, -OC(O)-(C<sub>1</sub>-C<sub>10</sub> alkylene)-NR<sup>6</sup>C(O)O-(alkylaryl), -P(O)(OR<sup>10</sup>)<sub>2</sub>, -(C<sub>1</sub>-C<sub>10</sub> alkylene)-OSi(alkyl)<sub>3</sub>, -CF<sub>3</sub>, -OCF<sub>3</sub>, halo, alkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyaryalkoxy, alkoxyiminoalkyl, alkyldioyl, allyloxy, aryl, arylalkyl, aryloxy, arylalkoxy, aroyl, aroyloxy, aroylaroyloxy, arylalkoxycarbonyl, benzoylbenzoyloxy, heteroaryl, heteroarylalkyl, heteroarylalkoxy, dioxolanyl, heterocyclyl, heterocyclylalkyl, heterocyclylcarbonyl, heterocyclylcarbonylalkoxy and -L-M;


wherein optionally one or more carbon atoms of the -(C<sub>01</sub>-C<sub>30</sub> alkylene)- radical of Q<sup>1</sup>, Q<sup>2</sup>, Q<sup>3</sup>, Q<sup>4</sup> and Q<sup>5</sup> is independently replaced by -O-, -C(O)-, -CH=CH-, -C≡C-, -N(alkyl)-, -N(alkylaryl)- or -NH-;

G is selected from the group consisting of a sugar residue, disugar residue, trisugar residue, tetrasugar residue, sugar acid, amino sugar, amino acid residue, oligopeptide residue

Reply under 37 CFR 1.116  
 Expedited Procedure  
 Technology Center 1624  
 Attorney Docket No. CV06025US01



comprising 2 to 9 amino acids, trialkylammoniumalkyl ~~salts~~ and  $-S(O)_2-OH$ , wherein optionally the sugar residue, disugar residue, trisugar residue, tetrasugar residue, sugar acid, amino sugar, amino acid residue or oligopeptide residue of G is substituted with  $-L-M$ ;

L is selected from the group consisting of






wherein Me is methyl;



M is selected from the group consisting of




Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01



Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01



and



$R^2$  and  $R^3$  can be the same or different and each is independently selected from the group consisting of hydrogen, alkyl and aryl;

$R^6$ ,  $R^7$  and  $R^8$  can be the same or different and each is independently selected from the group consisting of hydrogen, alkyl, aryl and arylalkyl; and

each  $R^9$  is independently alkyl, aryl or arylalkyl.

Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01

each  $R^{10}$  is independently H or alkyl;

q is 0 or 1;

r is 0 or 1;

m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3, 4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5;

x1 is 1 to 10;

x2 is 1 to 10;

x3 is 1 to 10;

x4 is 1 to 10;

x5 is 1 to 10;

x6 is 1 to 10; and

x7 is 1 to 10;

with the proviso that at least one of  $Q^1$ ,  $Q^2$ ,  $Q^3$ ,  $Q^4$  and  $Q^5$  is -L-M or the sugar residue, disugar residue, trisugar residue, tetrasugar residue, sugar acid, amino sugar, amino acid residue or oligopeptide residue of G is substituted with -L-M, and

wherein each of the above alkyl, alkenyl, alkynyl, alkylene, alkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyaryloxy, alkoxyiminoalkyl, alkyldioyl, allyloxy, aryl, arylalkyl, aryloxy, arylalkoxy, aroyl, aroyloxy, aroylaroyloxy, arylalkoxycarbonyl, benzoylbenzoyloxy, heteroaryl, heteroarylalkyl, heteroarylalkoxy, dioxolanyl, heterocyclyl, heterocyclylalkyl, heterocyclylcarbonyl, or heterocyclylcarbonylalkoxy groups, when present, is independently substituted or unsubstituted.

2. (Original) The compound according to claim 1, wherein m, n and r are each zero, q is 1, p is 2, and Z is  $-\text{CH}_2-$ .

3. (Original) The compound according to claim 1, wherein m, n and r are each zero, q is 1, p is 2, and Z is  $-\text{CH}_2-$ ,  $Q^1$  is  $-\text{OR}^6$ , wherein  $R^6$  is hydrogen and  $Q^5$  is fluorine.

{W0301621.1}

Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01

4. (Currently Amended) The compound according to claim 1, wherein  $R^2$  and  $R^3$  are each preferably hydrogen.

5. (Original) The compound according to claim 1, wherein  $Q^1$  and  $Q^2$  are each independently selected from the group consisting of  $-OR^6$ ,  $-O(CO)R^6$ ,  $-O(CO)OR^9$  and  $-O(CO)NR^6R^7$ .

6. (Original) The compound according to claim 1, wherein  $Q^4$  is halo or  $-OR^6$ .

7. (Original) The compound according to claim 1, wherein  $Q^1$  is  $-OR^6$  wherein  $R^6$  is H.


8. (Original) The compound according to claim 1, wherein  $Q^1$ ,  $Q^2$ ,  $Q^3$ ,  $Q^4$  or  $Q^5$  is  $-L-M$ .

9. (Currently Amended) The compound according to claim 1, wherein  $Q^1$ ,  $Q^2$ ,  $Q^3$ ,  $Q^4$  or  $Q^5$  is  $-G$  or  $-(C_1-C_{30} \text{ alkylene})-G$ .

10. (Withdrawn) The compound according to claim 1, wherein  $G$  is selected from the group consisting of:



Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01



wherein R, R<sup>a</sup> and R<sup>b</sup> can be the same or different and each is independently selected from the group consisting of H, -OH, halo, -NH<sub>2</sub>, azido, alkoxyalkoxy or -W-R<sup>30</sup>;

W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R<sup>31</sup>)-, -NH-C(O)-N(R<sup>31</sup>)- and -O-C(S)-N(R<sup>31</sup>)-;

R<sup>2a</sup> and R<sup>6a</sup> can be the same or different and each is independently selected from the group consisting of H, alkyl, acetyl, aryl and arylalkyl;

$R^{3a}$ ,  $R^{4a}$ ,  $R^{5a}$ ,  $R^{7a}$ ,  $R^{3b}$  and  $R^{4b}$  can be the same or different and each is independently selected from the group consisting of H, alkyl, acetyl, arylalkyl, -C(O)alkyl and -C(O)aryl;

$R^{30}$  is independently selected from the group consisting of  $R^{32}$ -substituted T,  $R^{32}$ -substituted-T-alkyl,  $R^{32}$ -substituted-alkenyl,  $R^{32}$ -substituted-alkyl,  $R^{32}$ -substituted-cycloalkyl and  $R^{32}$ -substituted-cycloalkylalkyl;

$R^{31}$  is independently selected from the group consisting of H and alkyl;


T is independently selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;

$R^{32}$  is 1 to 3 substituents which are each independently selected from the group consisting of H, halo, alkyl, -OH, phenoxy, -CF<sub>3</sub>, -NO<sub>2</sub>, alkoxy, methylenedioxy, oxo, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, -N(CH<sub>3</sub>)<sub>2</sub>, -C(O)-NHalkyl, -C(O)-N(alkyl)<sub>2</sub>, -C(O)-alkyl, -C(O)-


Reply under **37 CFR 1.116**  
 Expedited Procedure  
 Technology Center 1624  
 Attorney Docket No. CV06025US01

alkoxy and pyrrolidinylcarbonyl; or  $R^{32}$  is a covalent bond and  $R^{31}$ , the nitrogen to which it is attached and  $R^{32}$  form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolinyl or morpholinyl group, or a alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyl or morpholinyl group.

11. (Withdrawn) The compound according to claim 10, wherein G is selected from:




Reply under **37 CFR 1.116**  
 Expedited Procedure  
 Technology Center 1624  
 Attorney Docket No. CV06025US01




wherein Ac is acetyl and Ph is phenyl.

12. (Previously Presented) The compound according to claim 1, wherein optionally one or more carbon atoms of the  $-(C_1-C_{30} \text{ alkylene})-$  radical of  $Q^1$ ,  $Q^2$ ,  $Q^3$ ,  $Q^4$  and  $Q^5$  is independently replaced by  $-O-$ .


13. (Original) The compound according to claim 1, wherein L is



14. (Original) The compound according to claim 1, wherein M is




15 (Original) The compound according to claim 1, wherein M is




Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01


16. (Original) The compound according to claim 1, wherein M is



17. (Original) The compound according to claim 1, wherein M is




18. (Original) The compound according to claim 1, wherein M is




19. (Original) The compound according to claim 1, which is selected from the group consisting of


Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01




(II),



(III),



(IV),




(V),


{W0301631.1}

14


Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01



(VI),






(VII),




(VIII),


Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01




Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01



(XII),



(XIII), and



(XIV).

20. (Previously Presented) A pharmaceutical composition for the treatment of atherosclerosis, hypercholesterolemia, sitosterolemia, diabetes mellitus, obesity, stroke, lowering a concentration of cholesterol, phytosterol or 5 $\alpha$ -stanol in plasma of a mammal, treating demyelination or treating Alzheimer's disease and/or regulating levels of amyloid  $\beta$  peptides in a



Reply under 37 CFR 1.116  
Expedited Procedure  
Technology Center 1624  
Attorney Docket No. CV06025US01

subject comprising a therapeutically effective amount of a compound of claim 1 in a pharmaceutically acceptable carrier.

21. (Original) A pharmaceutical composition comprising a cholesterol-lowering effective amount of a compound of claim 1 in a pharmaceutically acceptable carrier.

22. (Previously Presented) A method of treating atherosclerosis, hypercholesterolemia, sitosterolemia, ~~diabetes mellitus, obesity, stroke,~~ lowering a concentration of cholesterol, phytosterol or 5 $\alpha$ -stanol in plasma of a mammal, ~~treating demyelination or treating Alzheimer's disease or regulating a level of an amyloid  $\beta$  peptide in a~~ subject comprising the step of administering to a subject in need of such treatment an effective amount of a compound of claim 1.

23. (Original) A method of lowering cholesterol level in plasma of a mammal in need of such treatment comprising administering a pharmaceutically effective amount of the compound of claim 1.